Provider Information

Insurance Coverage for Lymphedema Compression Garments and Supplies

Lymphedema Treatment Act and Who is Affected

Medicare now reimburses for compression garments and related supplies for individuals with lymphedema, following the passage of the Lymphedema Treatment Act in late 2022. This guide outlines key information specific to traditional Medicare coverage; however, many commercial and Medicaid plans often adopt similar policies, so it's recommended to verify reimbursement eligibility directly with each payer.



The official Medicare page for the new coverage can be found at go.cms.gov/48PxTKf.

• Daytime Garments

Ouantities Covered

- 3 sets (one garment for each affected body part) every six months.
- Nighttime Garments
 2 sets (one garment for each affected body part)
 every two years.
- Bandaging Supplies
 No set limit.
- Accessories
 No set limit, determined on a case-by-case basis depending on needs of the patient.

Compression Supplies Covered

- Custom and standard-fit daytime and nighttime garments.
- Custom and standard-fit gradient compression wraps with adjustable straps.
- Bandaging supplies for the initial and ongoing phases of treatment.
- Accessories including but not limited to lining, padding, zippers, donning and doffing aids.






Requirements for Coverage

- A lymphedema diagnosis, applicable codes are:
 - Q82.0 Hereditary lymphedema
 - **I89.0** Lymphedema, not elsewhere classified
 - **I97.2** Postmastectomy lymphedema syndrome
 - **197.89** Other postprocedural complications and disorders of the circulatory system, not elsewhere classified
- · A prescription for the item or items.
- Clinical notes must include the stage of lymphedema, and if a custom garment is ordered, notes must indicate why a standard-fit garment is not indicated for the patient.

Guidance on How to Diagnose Lymphedema

Lymphedema is a progressive disease marked by chronic swelling in one or more parts of the body. A lymphedema diagnosis is required by Medicare and other insurance providers in order for a patient to be eligible for coverage of compression garments and supplies. Providers can diagnose lymphedema 90% of the time by evaluating risk factors and conducting a physical examination. Imaging studies are usually not necessary. For in-depth information on diagnosis, staging, and management, see the three article links to the right.







bit.ly/LEdiagnosis2



bit.ly/LEdiagnosis3

Codes and Reimbursement Rates for Lymphedema Compression Supplies



The codes and corresponding reimbursement rates can be found in the CMS DMEPOS Fee Schedule.

Where can compression supplies be purchase?



Use our **Supplier Directory** to help locate suppliers who may meet your patient's needs. **lymphedemaproject.org/find-a-supplier/**

What details and clinical notes must be included with the prescription?

The prescription should include clear product descriptions that correspond to the new codes including the body part, type of garment, compression level if applicable, and whether a standard-fit or custom-fit item is needed. The clinical record must provide the rationale for the prescribed items and clearly document the medical necessity for each garment. The clinical notes, from prescribing physician or other medical provider must include a description of the lymphedema by body part, including stage/severity, followed by the compression garment type that is prescribed to treat that body part.

Example 1: Custom-fit (made to measure) daytime flat knit and nighttime padded garment request:

Insufficient: Bilateral lower extremity garments needed for lymphedema maintenance.

Sufficient: Patient presents with severe, bilateral, congenital primary lymphedema. Patient requires daytime, custom-fit, flat knit, close-toed, thigh-high garments for each leg. Custom-fit is necessary to accommodate and deliver gradient compression throughout the affected extremities due to their abnormal, disproportionate shape. Quantity 3 for each leg is needed for proper wash and wear. A nighttime custom-fit garment for each leg, toes to upper thigh, is also necessary to soften fibrotic tissue and prevent nighttime reaccumulation of swelling. Quantity 2 for each leg is needed for proper wash and wear.

Example 2: Standard-fit (ready to wear) daytime garment request:

Insufficient: Unilateral upper extremity compression garments required.

Sufficient: Patient presents with mild, unilateral, left, upper extremity lymphedema, secondary to tumor resection in 2020. Lymphedema extends throughout the extremity from fingers to the axilla. Patient requires a standard-fit, circular knit, compression sleeve and a separate compression glove. Quantity 3 of each item is needed for proper wash and wear.

